

DENDROCHRONOLOGY IN JAPAN

ENGLISH SUMMARY

RESEARCH REPORT OF
THE NARA NATIONAL CULTURAL
PROPERTIES RESEARCH INSTITUTE No. 48

NARA 1990

Contents

Preface

Explanatory notes

I. Dendrochronology and its Previous Researches	3
by TANAKA Migaku	
A. Outline of Dendrochronology	3
B. Previous Researches	7
1. Beginnings of dendrochronology and progress in Europe and North America	7
2. History of dendrochronology in Japan	11
II. Samples for Dendrochronological Research and its Methodology	18
by MITSUTANI Takumi	
A. Samples and Measurement of Width	18
1. Types of samples	18
2. Sample acquisition and preparation	19
3. Measurements of tree ring width	20
B. Cross-Correlation of Tree-Ring Patterns	21
1. Drawing tree-ring pattern graphs	21
2. Cross-correlation of tree-ring patterns by numerical data	23
3. Recognition of key signature patterns	27
III. Examination of Ring Patterns Utilizing Living Trees	29
by MITSUTANI Takumi	
A. Examination of Features of Ring Patterns of Living <i>Hinoki</i> Cypress	29
1. Acquisition of <i>hinoki</i> cypress samples	29
2. Ring patterns of the same tree taken radially in different directions at the same level	31
3. Ring patterns at the center and near the exterior of the same tree at the same level	34
4. Ring patterns taken at different height of the same tree	34
5. Ring patterns taken from different trees in the same	

region	38
6. Ring patterns taken from trees of different region	45
B. Ring Patterns of Trees of Non-Cypress Species	49
1. Ring patterns of <i>hiba</i> arborvitae and <i>sugi</i> cedar	50
2. Ring patterns of <i>kōyamaki</i> pine and other species	55
IV. Building Index Master Chronology	60
by MITSUTANI Takumi	
A. Building Standard Chronology of <i>Hinoki</i> Cypress	60
1. Building standard chronology of modern trees	60
2. Confirmation and extensions of standard chronology	67
a. Standard chronology derived from structural parts of the Prayers Hall of the Nigatsu-dō 二月堂 at Tōdai-ji 東大寺, Nara Prefecture	68
b. Standard chronology derived from artifacts discovered at the site of the Kiyosu 清洲 Castle Town, Aichi Prefecture	71
c. Standard chronology derived from artifacts discovered at the Kusado Sengen 草戸千軒 site, Hiroshima Prefecture	73
d. Standard chronology derived from artifacts discovered at the Toba 鳥羽 Detached Palace site, Kyoto Prefecture	74
e. Standard chronology derived from artifacts discovered in the Heijō 平城 Capital, Nara Prefecture	76
f. Extension of standard chronology beyond the Time of Christ	78
B. Standard Chronology Derived from Non-Cypress Species	79
1. Comparison between ring patterns derived from different species of trees in the same region	79
2. Comparison between ring patterns derived from different species of trees in different regions	80
3. Standard chronology of <i>sugi</i> cedar	82
a. Standard chronology derived from living <i>sugi</i> cedar	82
b. Standard chronology derived from archaeologically discovered artifacts and structural parts in the Tohoku area	83
c. Extension of the standard chronology of <i>sugi</i> in the Tohoku area	86
d. Mean ring pattern of securely dated wooden artifacts dis- covered in Shizuoka Prefecture	87
4. Standard chronology of <i>kōyamaki</i> pine	89
a. Standard chronology derived from artifacts discovered in Nara Prefecture	89

b. Floating chronologies	92
V. Application of Index Master Chronology	94
by MITSUTANI Takumi and TANAKA Migaku	
A. Introduction	94
B. Application of Dendrochronology	98
1. Dendrochronological dates of living trees	98
2. Dendrochronological Dates of Artifacts ¹	99
a. Frames of a well: Ochiai 落合Ⅲ site, Iwate Prefecture	99
b. Posts: Hotta no Saku 弘田柵 fort site*, Akita Prefecture	99
c. Structural parts of houses buried at the Kurumidate 胡桃館 site, Akita Prefecture	100
d. Posts: Kinowa no Saku 城輪柵 fort site*, Yamagata Prefecture	101
e. Coffin: Nanamawari-Kagamizuka Kofun 七廻り鏡塚古墳 burial site, Tochigi Prefecture	101
f. Funerary urn: site of the Castle Town of the Asakura 朝倉 Family of Ichijōdani 一乗谷**, Fukui Prefecture	102
g. Wooden objects: Yamagi 山木 site, Shizuoka Prefecture	102
h. Preserved pillars: Shida Gunga 志太郡衙 county-office site*, Shizuoka Prefecture	103
i. Wooden objects: Kiyosu 清洲 Castle Town site, Aichi Prefecture	104
j. Structural parts: a bridge attributed to the Seta no Karahashi 瀬田唐橋, Shiga Prefecture	104
k. Preserved pillars: Miyamachi 宮町 site, Shiga Prefecture	104
l. Coffin: Jōdoji Kofun 浄土寺古墳 burial site, Kyoto Prefecture	106
m. Coffin: Kawaradani 瓦谷 site, Kyoto Prefecture	107
n. Parasol-shaped object: Moat of the Kondayama Kofun 誉田山古墳 burial site (attributed to the Mausoleum of Emperor Ōjin 応神), Osaka Prefecture	107
o. Coffin: Kyōzuka Kofun 経塚古墳 burial site, Osaka Prefecture	107
p. Frames of a well: Ori'ono 遠里小野 site, Osaka Prefecture	108

1 In this section, * and ** respectively denote a "Historic Site" and a "Historic Site of Special Significance" designated by the national government.

- q. Parasol-shaped objects: Shijō Kofun 四条古墳 burial site, Nara Prefecture 108
 - r. Parasol-shaped objects: Obaka Kofun 小墓古墳 burial site, Nara Prefecture 109
 - s. Part of a water clock: Mizu'ochi 水落 site*, Nara Prefecture 109
 - t. Preserved lower portions of pillars: Lower stratum of the Hokke-ji 法華寺 temple site, Nara Prefecture 110
 - u. Votive tablet of a horse: Higashi-Ni Bō 東二坊[East Second Ward] on the Nijō Ōji 二条大路[Second Major Street], Heijō 平城 Capital, Nara Prefecture 111
 - v. Frames of a well: 13th and 14th Tsubo 坪[Blocks], Ichī Bō 一坊[First Ward] on Hachijō Ōji 八条大路[Eighth Major Street], Ukyō 右京[Western Sector], Heijō Capital 112
 - w. Trough: location of the historically known Masuda no Ike 益田池 pond, Nara Prefecture 112
 - x. Artifacts discovered at the Kusado Sengen 草戸千軒 site, Hiroshima Prefecture 112
 - y. Bases of a circular wooden-bent boxes: Shimokawazu 下川津 site, Kagawa Prefecture 113
 - z. Lower preserved portion of a pillar: Ōno 大野 Castle site**, Fukuoka Prefecture 114
3. Dendrochronological dates of architectural structures²...114
- a. Boards used for a wall: Kagura-den 神楽殿 [Hall of Ritual Music and Dance]*, Ha'ushiwake 波宇志別 Shrine, Akita Prefecture 114
 - b. Pillar: Main Hall*, Wakamiya Hachiman 若宮八幡 Shrine, Nagano Prefecture 115
 - c. Structural parts: Three story pagoda*, Kiyomizu-dera 清水寺 temple, Kyoto Prefecture 116
 - d. Structural parts: Main Hall*, Gaya-in 伽耶院 temple, Hyogo Prefecture 116
 - e. Central pillar: Five story pagoda**, Hōryū-ji 法隆寺 temple, Nara Prefecture 116
 - f. Central pillar: Three story pagoda**, Hokki-ji 法起寺 temple, Nara Prefecture 118

² In sections 3 and 4,* indicates an "Important Cultural Property,"

**a "National Treasure."

g.	Structural parts: Ōtataneko 大直禰子 Shrine*, a subordinate shrine of the Ōmiwa 大神 Shrine, Nara Prefecture	118
h.	Structural parts: Main Hall*, Hōtō-ji 宝幢寺 temple, Nara Prefecture	118
4.	Dendrochronological dates of art objects	120
a.	Lacquered wooden bent-box kept at a temple in Kyoto Prefecture	120
b.	Seat of the Bhaisjyaguruviduryaprabha statue** in the Image Hall, Hōryū-ji, Nara Prefecture	120
c.	Hyakuman - tō 百万塔 [one million miniature pagodas] at Hōryū-ji, Nara Prefecture	120
d.	Guardian figure (the one in a pair who keeps the mouth colsed) in the South Gate, Tōdai-ji 東大寺 temple, Nara Prefecture	122
e.	Seated Amitabha at Ganki-ji 岩崎寺 temple, Yamaguchi Prefecture	122
f.	Seated Amitabha at Hōkō-ji 法光寺 temple, Yamaguchi Prefecture	124
g.	Standing Arya-avalokitesvara at Hōkō-ji temple, Yamaguchi Prefecture	124
h.	Standing Vaisravana at Hōkō-ji temple, Yamaguchi Prefecture	124
i.	Standing Acalanatha at Hōkō-ji temple, Yamaguchi Prefecture	125
j.	Standing Virupaksa at Gatsurin-ji 月輪寺 temple, Yamaguchi Prefecture	125
5.	Dendrochronological dates of buried forests	126
a.	<i>Sugi</i> cedar in the Yamamoto Country, Akita Prefecture	126
b.	<i>Sugi</i> cedar in the Yuri County, Akita Prefecture	126
c.	<i>Sugi</i> cedar in the Mogami Country, Yamagata Prefecture	127
d.	<i>Hinoki</i> cypress in Susono City, Shizuoka Prefecture	127
VI.	Dendroclimatology and its Applications	128
	by SATŌ Tadanobu and YAJIMA Atsushi	
A.	Detecting Characteristic Long - Term Climatic Fluctuations Based on Changes in the Tree-Ring Width	128
1.	Reconstruction of palaeo-climate and the examination of data	128
2.	Standardization of tree-ring width data	130
a.	Method of standardization	130

b. Tree-ring width data of living trees 131	
3. Climatic data	133
B. Detecting Characteristic Climatic Fluctuations	140
1. Correlations between tree-ring width and climatic factors	140
2. Estimation of climatic data utilizing correlation functions	144
a. Overview of the methods of estimation 144	
b. Results of the estimation 145	
3. Estimation of climatic data by the Kalman filtering algorithm	147
C. Conclusions	157
VII. Development of Dendrochronology in Japan	159
by TANAKA Migaku	
References	163
English summary	169

List of Figures

I — 1	Building the Index Master Chronology	6
I — 2	Index Master Chronology used in other parts of the world (as of 1987), after Schweingruber, 1988	10
II — 1	Core specimen extractor and sample-support	19
II — 2	Increment borer and sample-support	19
II — 3	Incremental measuring machine	20
II — 4	Cross-dating by skeleton plots	21
II — 5	Tree-ring patterns: standardized (thick line) and measured	23
II — 6	Key-signature rings (thick lines) and the detection method	28
III — 1	Locality where samples of living <i>hinoki</i> cypress were collected	30
III — 2	Ring patterns taken radially in three directions from at the same level of a tree	32
III — 3	Comparison of ring patterns taken toward the exterior of a tree (rings formed in tree's late life)	35
III — 4	Comparison of ring patterns taken in the center of a tree (rings formed in tree's early life)	35
III — 5	Ring patterns of Samples No. 2 and No. 14 from Agematsu 上松, Nagano Prefecture	40
III — 6	Ring patterns of a sample from Agematsu and one from Tsukechi 付知, Gifu Prefecture	46
III — 7	Relationship between distance of two localities where samples were collected and t value [horizontal axis : distance ; vertical axis : t values]	47
III — 8	Ring patterns according to different localities	48
IV — 1	Standard chronology of a living tree: between 1009 and 1100	62
IV — 2	Standard chronology of a living tree: between 1100 and 1400	63
IV — 3	Standard chronology of a living tree: between 1400 and 1700	64
IV — 4	Standard chronology of a living tree: between 1700 and 1984	65
IV — 5	Key signature rings in Standard Chronology A of living trees	66
V — 1	Where three different samples were taken on the cross-section of the same tree	95
V — 2	Standard chronology derived from six samples of <i>hinoki</i> cypress	97
V — 3	Basal plate of a funerary urn : site of Castle Town of Asakura Family of Ichijōdani, Fukui Prefecture	103
V — 4	Preserved lower portion of a pillar : Miyamachi site, Shiga Prefecture	105

V—5	Votive table of a horse: Higashi-Ni Bō on the Nijō Ōji, Heijō Capital, Nara Prefecture	111
V—6	Main building: Wakamiya Hachiman Shrine, Nagano Prefecture...	115
V—7	Cross-section of the central pillar: Five story pagoda, Hōryū-ji temple, Nara Prefecture	117
V—8	Overhaul of the Main Hall: Ōtataneko Shrine, Nara Prefecture...	119
V—9	Lacquered wooden-bent box of a temple in Kyoto Prefecture.....	121
V—10	One million miniature Pagodas of Hōryū-ji temple, Nara Prefecture	121
V—11	Guardian figure in the South Gate, Todai-ji temple; Nara Prefecture and its inner structure	123
V—12	Seated Amitabha at Hōkō-ji temple, and Standing Virupaksa at Gatsurin-ji temple, Yamaguchi Prefecture	125
VI—1	Comparison of ring width spectrum over years in Lapland (1463–1960)[vertical axis: ring width spectrum; horizontal axis: years] (upper) and mean temperature (from April to September) spectrum over years (1659–1973)[vertical axis: mean temperature spectrum; horizontal axis: year](lower)	129
VI—2	Moving averaged ring pattern (ten year mean) of <i>hinoki</i> cypress in the Kiso region [vertical axis: ring width; horizontal axis: year]	131
VI—3	Drastic changes in tree ring width which took place around 1840 [vertical axis: gap in millimeters; horizontal axis: year]	131
VI—4	Spectrum of tree ring width to which the Loess filtering has been applied [horizontal axis: year]	132
VI—5	Fluctuation in tree ring width: measured (upper) and standardized after correcting base line fluctuation (lower)[vertical axis: ring width in millimeters; horizontal axis: year]	132
VI—6	Fluctuations in ring width of <i>hinoki</i> cypress year by year: measured [vertical axis: ring width in millimeters; horizontal axis: year]	134
VI—7	Fluctuation in ring width of <i>hinoki</i> cypress: standardized by smoothing of the spline algorithm [vertical axis: ring width in millimeters; horizontal axis: year]	135
VI—8	Localities in Nagano Prefecture where per annum rainy day ³ data were obtained	137
VI—9	Descriptions of weather in the <i>Bokuō Nikki</i> 墨翁日記 (parentheses indicate the applicable descriptions.)	137
VI—10	Comparison between moving averaged per annum rainy days (ten year mean)[vertical axis: per annum rainy days; horizontal axis: year](upper) and moving averaged fluctuation in precipitation (ten	

3 In this chapter, by “per annum rainy days” we mean the total number of rainy days from April to September in each year.

	year mean) in Kiso-Fukushima [vertical axis: precipitation in millimeters; horizontal axis: year](lower)	137
VI—11	Power spectra of ring width (in millimeters) of <i>hinoki</i> cypress in the Kiso region (upper) of mean temperature (in degrees in centigrade) in Nagano City (middle), and of per annum rainy days (in days) in Nagano City (lower)[vertical axis: power spectrum; horizontal axis: year]	140–141
VI—12	Cross correlation coefficient of per annum rainy days and tree ring width (upper) and cross correlation coefficient of mean temperature and tree ring width (lower) [vertical axis: cross correlation coefficient; horizontal axis: year]	142
VI—13	Fluctuation in mean temperature in England [vertical axis: temperature in degrees in centigrade; horizontal axis: year](upper) and fluctuation in frequency of floods of Kamo 鴨, River in Kyoto Basin for fifty years [vertical axis: number of floods; horizontal axis: year]	143
VI—14	Comparison of per annum rainy days estimated by cross correlation coefficient and actual data of rainy days [solid line: estimated; broken line: actual; vertical axis: rainy days; horizontal axis: year]	147
VI—15	Power spectra of estimated per annum rainy days and of actual data of rainy days [solid line: estimated; broken line: actual; vertical axis: power spectra; horizontal axis: year]	147
VI—16	Patterns of per annum rainy days estimated from tree-ring width by cross correlation coefficient and of actual data of rainy days [solid line: estimated; broken line: actual; vertical axis: rainy days; horizontal axis: year]	147
VI—17	Process of the identification of system parameters by autoregressive moving average coefficients of Kalman filtering algorithm [vertical axis: system parameter; horizontal axis: iteration]	151–153
VI—18	Fluctuation in per annum rainy days estimated from tree ring data [vertical axis: rainy days; horizontal axis: year]	154
VI—19	Power spectra of estimated per annum rainy days over years and of actual per annum rainy days [vertical axis: power spectrum; horizontal axis: year]	155
VI—20	Fluctuation of estimated per annum rainy days for one thousand years [vertical axis: rainy days; horizontal axis: year]	156
VII—1	Ring patterns taken from preserved lower portions of four pillars discovered at the Mawaki 真脇 site of the Jomon Period, Ishikawa Prefecture	160
VII—2	Ring patterns taken from three pieces of timber discovered at the Paleolithic Tomizawa 冨沢 site, Miyagi Prefecture	161

List of Tables

II — 1	t values for two pairs of the cross-correlations of tree-ring patterns	26
III — 1	Localities, where samples of living <i>hinoki</i> cypress trees were collected	30
III — 2	Correlation t values of ring patterns taken radially in three directions from the same sample [A1 — A3: sample numbers; outside of the parenthesis: t value; number in the parenthesis: correlation coefficient]	32
III — 3	Comparison of ring patterns taken in the center of a tree (rings formed in tree's early life) and taken toward the exterior of a tree (rings formed in tree's late life) [ED1: ring pattern showing tree's late life (last 100 years); ED2: ring pattern of tree's early life (from the center to ED1); bold characters: t values above 3.5]	33
III — 4	Correlation t values among samples taken radially in four directions at two levels of the same tree [A1—B2: sample number; number in parenthesis: level at which each sample was taken]	36
III — 5	Correlation t values among ring patterns taken at different levels of the same tree	37
III — 6	Correlation t values among ring patterns taken at different levels and the average ring pattern of samples taken at Agematsu, Nagano Prefecture	38
III — 7	Correlation t values among the ring patterns of twenty samples taken at Agematsu, Nagano prefecture	40
III — 8	Correlation t values among eighteen ring patterns taken at Miura 三浦, Nagano Prefecture	41
III — 9	Correlation t values among six ring patterns taken at Kōrigase 氷ヶ瀬, Nagano Prefecture	44
III — 10	Correlation t values among five ring patterns taken at Komata 小俣, Nagano Prefecture	44
III — 11	Correlation t values among seventeen ring patterns taken at Tsukechi, Gifu Prefecture	42
III — 12	Correlation t values among eighteen ring patterns from 150 samples taken at Kosaka-Ōbora 小坂大洞, Gifu Prefecture	43
III — 13	Correlation t values among eleven ring patterns from 211 samples taken at Kosaka-Ōbora, Gifu Prefecture	44
III — 14	Correlation t values among six ring patterns taken at Owase 尾鷲, Mie Prefecture	44

III—15	Correlation t values among two ring patterns taken at Kōyasan 高野山, Wakayama Prefecture	44
III—16	Correlation t values among six ring patterns taken at Yanase 魚梁瀬, Kochi Prefecture	44
III—17	Correlation t values among the average ring patterns of the ten localities above	46
III—18	Correlation t values between the average ring patterns of the Tsukechi samples and of the Yanase samples	48
III—19	Non-cypress species and the localities where samples were collected	49
III—20	Correlation t values among ten ring patterns of <i>hiba</i> arborvitae taken at Ōhata 大畑, Aomori Prefecture	50
III—21	Correlation t values among ten ring patterns of <i>hiba</i> arborvitae taken at Kawauchi 川内, Aomori Prefecture	51
III—22	Correlation t values among ten ring patterns of <i>hiba</i> arborvitae taken at Yokohama 横浜, Aomori Prefecture	51
III—23	Correlation t values among eight ring patterns of <i>hiba</i> arborvitae taken at Masukawa 増川, Aomori Prefecture	51
III—24	Correlation t values among ten ring patterns of <i>hiba</i> arborvitae taken at Imabetsu 今別, Aomori Prefecture	52
III—25	Correlation t values among eight ring patterns of <i>hiba</i> arborvitae taken at Kanagi 金木, Aomori Prefecture	52
III—26	Correlation t values among seven ring patterns of <i>hiba</i> arborvitae taken at Kawai 川井, Iwate Prefecture	52
III—27	Correlation t values among the average ring patterns of <i>hiba</i> arborvitae taken at different localities	53
III—28	Correlation t values among ten ring patterns of <i>sugi</i> cedar taken at Fujisato 藤里, Akita Prefecture	55
III—29	Correlation t values among ten ring patterns of <i>sugi</i> cedar taken at Akita 秋田, Akita Prefecture	56
III—30	Correlation t values among nine ring patterns of <i>sugi</i> cedar taken at Yanase, Kochi Prefecture	56
III—31	Correlation t values among four ring patterns of <i>sugi</i> cedar taken in the Yaku 屋久 Island, Kagoshima Prefecture	56
III—32	Correlation t values among the mean ring patterns of <i>sugi</i> cedar taken at the four localities above	56
III—33	Correlation t values among eight ring patterns of <i>sawara</i> cypress at Ōtaki 王滝, Nagano Prefecture	57
III—34	Correlation t values among four ring patterns of <i>kōyamaki</i> pine at Ōtaki, Nagano Prefecture	58
III—35	Correlation t values among three ring patterns of <i>mizunara</i> oak at Kawauchi, Aomori Prefecture	58
III—36	Correlation t values among three ring patterns of <i>mizunara</i> oak at	

	Ottomo 乙供, Aomori Prefecture	58
III-37	Correlation t values among ring patterns of <i>buna</i> beech taken at three localities	59
IV-1	Correlation t values among ring patterns of Group I of lumber samples taken from the Prayers Hall of the Nigatsu-dō at Tōdai-ji, Nara Prefecture	70
IV-2	Correlation t values among ring patterns of Group II of lumber samples taken from the Prayers Hall of the Nigatsu-dō at Tōdai-ji, Nara Prefecture	71
IV-3	Correlation t values among ring patterns of Group III of lumber samples taken from the Prayers Hall of the Nigatsu-dō at Tōdai-ji, Nara Prefecture	71
IV-4	Key signature rings of Standard Chronology B derived from lumber of the Prayers Hall of the Nigatsu-dō at Tōdai-ji, Nara Prefecture	71
IV-5	Correlation t values among ring patterns of artifacts discovered at the site of the Kiyosu Castle Town, Aichi Prefecture [* indicates non-correlation at that point.]	72
IV-6	Key signature rings of Standard Chronology C derived from artifacts discovered at the site of the Kiyosu Castle Town, Aichi Prefecture	72
IV-7	Correlation t values among ring patterns of five nose rings discovered at the Kusado Sengen site, Hiroshima Prefecture [* indicates non-correlation at that point.]	73
IV-8	Correlation t values among ring patterns of twelve artifacts discovered at the Kusado Sengen site, Hiroshima Prefecture [* indicates non-correlation at that point.]	75
IV-9	Key signature rings of the mean ring pattern of samples from the Kusado Sengen site, Hiroshima Prefecture	75
IV-10	Correlation t values among ring patterns of six artifacts discovered at the Toba Detached Palace site, Kyoto [* indicates non-correlation at that point.]	75
IV-11	Key signature rings in Standard Chronology D	76
IV-12	Key signature rings in Standard Chronology E derived from artifacts discovered in the Heijō Capital	76
IV-13	Correlation t values among ring patterns of six artifacts discovered in the Heijō Capital, Nara Prefecture [* indicates non-correlation at that point.]	77
IV-14	Correlation t values among ring patterns derived from artifacts discovered in the Yayoi and Kofun Periods sites [* indicates non-correlation at that point.]	78
IV-15	Correlation t values among ring patterns of six different species grown in Nagano Prefecture	81
IV-16	Correlation t values among ring patterns of four different species	

	grown in Aomori Prefecture	81
IV-17	Correlation t values among ring patterns of different species grown in different regions	81
IV-18	Correlation t values among ring patterns taken from artifacts discovered in Yamagata Prefecture [* indicates non-correlation at that point.]	84
IV-19	Correlation t values among ring patterns taken from structural parts of architecture in Akita Prefecture [* indicates non-correlation at that point.]	85
IV-20	Correlation t values among ring patterns taken from frames of wells in Iwate Prefecture [* indicates non-correlation at that point.]	85
IV-21	Correlation t values among ring patterns taken from samples excavated at the Hotta no Saku fort site, Yamagata Prefecture [* indicates non-correlation at that point.]	88
IV-22	Correlation t values among ring patterns taken from artifacts discovered in Shizuoka Prefecture [* indicates non-correlation at that point.]	88
IV-23	Correlation t values among ring patterns taken from preserved lower portions of pillars excavated in the Heijō Palace site and Hokke-ji temple site, Nara Prefecture [* indicates non-correlation at that point.]	90
IV-24	Correlation t values among ring patterns taken from artifacts discovered at the Shijō Kofun burial site, Nara Prefecture [* indicates non-correlation at that point.]	93
IV-25	Correlation t values among ring patterns taken from a coffin discovered at the Kariya 雁屋 site, Osaka Prefecture [* indicates non-correlation at that point.]	93
V-1	Dendrochronological date: Ochiai III site, Iwate Prefecture	100
V-2	Dendrochronological date: Hotta no Saku fort site, Akita Prefecture	100
V-3	Dendrochronological date: Kurumidate site, Akita Prefecture	100
V-4	Dendrochronological date: Yamagi site, Shizuoka Prefecture	103
V-5	Dendrochronological date: Shida Gunga county office site, Shizuoka Prefecture	103
V-6	Dendrochronological date: Kiyosu Castle Town site, Aichi Prefecture	105
V-7	Dendrochronological date: Bent of the ancient Seta no Karahashi bridge, Shiga Prefecture	105
V-8	Dendrochronological date: Miyamachi site, Shiga Prefecture	105
V-9	Dendrochronological date: Shijō Kofun burial site, Nara Prefecture	109
V-10	Dendrochronological date: Obaka Kofun burial site, Nara Prefecture	

	109
V-11 Dendrochronological date: Lower stratum of the Hokke-ji temple site, Nara Prefecture	110
V-12 Dendrochronological date: Kusado Sengen site, Hiroshima Prefecture	113
V-13 Dendrochronological date: Shimokawazu site, Kagawa Prefecture	113
V-14 Dendrochronological date: Hau'shiwake Shrine, Akita Prefecture	115
V-15 Dendrochronological date: Gaya-in temple, Hyogo Prefecture	116
V-16 Dendrochronological date: Ōtataneko Shrine, Nara Prefecture	119
V-17 Dendrochronological date: Hōtō-ji temple, Nara Prefecture	119
V-18 Dendrochronological date: Lacquered wooden bent-box at a temple in Kyoto Prefecture	121
V-19 Dendrochronological date: Hyakuman - tō [one million miniature pagodas] at Hōryū-ji, Nara Prefecture	121
V-20 Dendrochronological date: Guardian figure in the South Gate, Tōdai-ji temple, Nara Prefecture	123
V-21 Dendrochronological date: Seated Amitabha at Ganki-ji temple, Yamaguchi Prefecture	123
V-22 Dendrochronological date: Seated Amitabha at Hōkō-ji temple, Yamaguchi Prefecture	124
VI-1 Number of rainy days from April to September every year between 1813 and 1982 [left column: year; center column: rainy days between April and September; right column: sources]	138-139
VI-2 Square errors between estimated number of rainy days and the observed number (determination of degrees of p and q) [left: number of q ; center: number of p ; right: number of n]	146
VI-3 Square errors between estimated number of rainy days and the observed number (determination of degrees of p and q) [left: number of q ; center: number of p ; right: number of n]	150

Dendrochronology in Japan

English Summary

This report summarizes a research into dendrochronology and dendroclimatology conducted principally by archaeologist TANAKA Migaku 田中琢 and archeo-botanist MITSUTANI Takumi 光谷拓実 at the Nara National Cultural Properties Research Institute. The principal investigators are joined by SATŌ Tadanobu 佐藤忠信 of Disaster Prevention Research Institute at Kyoto University in the application of dendroclimatology. Our five year project initiated in 1985 has been supported by the Grant-in-Aid for Scientific Research program of the Japanese Ministry of Education. Prior to the presentation of our results, we first review previous research into the use of tree rings in Japan. We will proceed to discuss our studies, first on dendrochronology, and then on dendroclimatology.

Studies based on tree rings in Japan originally began with the aim of reconstructing palaeoclimate and the cycle of palaeoclimatic fluctuations. These measurements were calculated on the basis of the width of tree rings. However, such researches always utilized only one tree; the scholars did not attempt to cross-date more than one sample. This was the case in all pre-World War II studies.

The earliest example of dendroclimatological scholarship in Japan was HIRANO Ressuke's 平野烈介 research published in 1921. He calculated the total amount of growth of a *sugi* cedar based on the area of the cross-section of the tree. He observed fluctuations in the amount of yearly growth and determined that: "In the life of this particular Japanese cedar tree, we have clear evidence of a thirty-three year cycle." He identified this cycle as Brückner's cycle in climatology. Such studies were continued intermittently, and other hypothetical periodic cycles of long-term climatic fluctuation were proposed — 110 years, 350 years, and 700 years. The most famous research was by YAMAZAWA Kingorō 山沢金五郎, then the director of the Weather Station at Takayama, Gifu Prefecture. In 1930, he published the result of his measurements of the width of 802 rings of a Japanese cypress tree (*Chamaecyparis obtusa* Endl.), and he attributed these 802 rings to the years between 1119 to 1920. In our re-examination, however, we discovered that these rings were formed between 1118 and 1919.

Among the pre-World War II scholars interested in dendrochronology, architectural historian SEKINO Masaru 関野克 was an exception in that he not only followed the progress in this field in Europe and North America, especially the achievements of A.E. Douglass who could indeed cross-date

some floating chronologies, but also attempted to apply Douglass' work to Japanese trees. In 1943, Sekino measured the width of rings of structural parts of an eighth century building in Japan, and some of the ring patterns he drew still survive. Probably because of the social confusion and disturbances during and soon after World War II, however, Sekino did not make any further progress in his research.

Another epochal study was published by SHIDEI Tsunahide 四手井綱英 and his collaborators in 1944. Their project, conducted in Akita Prefecture, was methodologically significant because they utilized measurements from several hundred Japanese cedars rather than just one tree. They concluded that the fluctuation in the tree growth between 1928 and 1942, as evidenced in the different width of tree rings, was related to the precipitation coefficient¹, the growth of Japanese cedar became poor whenever the precipitation coefficient went above or below certain values.

The majority of postwar tree ring studies in Japan were climatological. Scholars became methodologically precise and were no longer satisfied with simple comparisons between ring patterns and climatic fluctuations. Nonetheless, meticulous data-gathering such as characterized Shidei's work was not yet common.

Two studies conducted during that period were worthy of note. One was TAKAHASHI Hiroaki's 高橋宏明 study. Following Shidei's example, Takahashi explored the relationship between tree growth and climatic factors, especially precipitation coefficient. He showed the correlation among the average width of rings of several trees at one specific site and the average width taken at different sites. The fact that trees of different ages and at different localities showed the same growth pattern indicated the connection with a common environmental factor—climate. This was the starting point of his research. Unfortunately, no scholars continued this line of research and thinking.

Another interesting study was conducted by NISHIOKA Hideo 西岡秀雄, who set out to date an architectural structure using dendrochronology. He measured the width of approximately 250 rings of the central pillar of the pagoda at Hōryū-ji 法隆寺, Nara Prefecture, the oldest surviving wooden structure in the world. He then compared the measurements with a ring pattern taken from a part of a different structure built in the 730s also at Hōryū-ji and concluded that the tree used in making the Hōryū-ji pagoda pillar was cut down before 607. Nishioka's work was significant because it

1 The ratio between amount of annual precipitation and annual average temperature

was the first use of tree-rings for the purpose of dating. Nonetheless, both tree samples from which he took ring patterns lacked the outer most part of the bark, which would have given precise dates of trees cut down. Consequently, Nishioka's work has been regarded by later scholars as hasty and imprecise and has been duly ignored.

Part of Japanese negative attitude toward dendrochronology came from scholars' belief in the impossibility of its application to Japanese trees. They believed without any testing that the climate and topography in Japan were so different from Arizona where Douglass developed dendrochronology that what was possible in Arizona would not be possible in Japan.

What tempted us to pursue dendrochronology at the Nara National Cultural Properties Research Institute was that we had excavated numerous wooden artifacts at archaeological sites. We had also conducted ethnographical/historical field research into a number of ancient temples, shrines, and other old wooden structures. Our knowledge about European and American achievements in this field also stimulated our research.

In 1970, one of our colleagues started to measure tree ring width. After he processed some samples, he reached the conclusion that it would be very difficult to apply dendrochronological methods to Japanese cypresses. Since most buildings in Japan were traditionally made from cypresses, the project was halted.

In 1979 we started a preliminary research and found the application of dendrochronology and dendroclimatology to Japanese trees possible. In 1985, our major studies in these two fields began, and we confirmed the possibility of the application of dendrochronology to archaeology, history, architectural history, and art history. We also made considerable progress in dendroclimatology. At the same time of our research, other dendrochronological studies were published. Our studies could be distinguished by the fact of having established a significantly long-term ring pattern.

The method for cross-correlation of tree-ring patterns that we used was basically same as that of Europe. In order to standardize the varying tree ring width, we used the five running mean method. To determine statistical significance, we used Student's "*t*" test. With sixty or more degrees of freedom, we compared the obtained *t* value with the 0.1 per cent significance level of *t*, which was 3.5. Correlations of ring patterns which grew over the same span of years normally produced *t* values greater than $t = 3.5$. Our experience showed that samples in a pair which yielded a *t* value greater than 3.5 did not necessarily cross-correlate. We always double-checked with the results of visual matching of pattern graphs with reference

to key signature patterns. The statistical calculations were simply a means of cross-correlating ring patterns.

The materials were taken from living forest trees, old structures, and wooden artifacts discovered at sites. The kinds of trees used in our studies included: a *hinoki* cypress *Chamaecyparis obtusa* Endl., a *sawara* cypress *Chamaecyparis pisifera* Endl., an *asunaro* arborvitae *Thujopsis dolabrata* Sieb. et Zucc., a *hiba* arborvitae *Thujopsis dolabrata* Sieb. et Zucc. var. *Hondai* Makino, a *kurobe* arborvitae *Thuja Standishii* Carr., a *tsuga* hemlock *Tsuga sieboldii* Carr., a *sugi* cedar *Cyptomeria japonica* D. Don, and a *kōyamaki* pine *Sciadopitys verticillata* Sieb. et Zucc. all of which were coniferous trees, as well as some deciduous trees, such as a *mizunara* oak *Quercus mongolica* Fischer ex T. var. *grosseserrata* (Bl.) Rehd. et Wils. and a *buna* beech *Fagus crenata* Bl..

The first step of our dendrochronological studies was to confirm the applicability of dendrochronology to Japanese trees and to determine an appropriate species for our studies. For the first purpose, we selected a Japanese cypress because it was native to Japan, and natural forest of this tree was distributed from the southern half of the Honshu island to southern Kyushu. Cypressess had also been most widely used for Japanese architecture, and were often discovered at archaeological sites. Utilizing modern Japanese cypress trees, we recognized the following features important for our studies: 1) ring patterns taken radially from the same tree tended to highly correlate with each other; 2) rings formed during trees' early life tended to show features unique to individual trees, this leading to the intelligence that for dendrochronological studies we should avoid rings near the center of a tree; 3) rings near or at the base of a tree often showed anomalies because of the expansion of the root; 4) given a certain region, rings of different trees showed very high correlations to one another. We could even cross-correlate samples taken from trees approximately 400 kilometers apart.

For the second goal, we tested a *kōyamaki* pine, a *sawara* cypress, an *asunaro* arborvitae, a *hiba* arborvitae, a *kurobe* arborvitae, a *tsuga* hemlock, a *sugi* cedar, a *mizunara* oak, related species of which were widely used in European dendrochronological studies, and a *buna* beech which was the most representative deciduous trees in Japan. Among these, we found the first seven species suitable for our studies. It is ironic that the last two species were not found to be applicable.

The next step was to build an index master chronology, which was the foundation of any dendrochronological studies. Our index master chronology of *hinoki* cypress trees spanned from 317, B.C. to 1984. It was

derived from: Standard Chronology A(1695-1983) taken from a modern tree; Standard Chronology B (1027-1755) taken from a structural part of a sixteenth century building at Tōdai-ji 東大寺, Nara; Standard Chronology C (751 - 1591) derived from artifacts discovered in the Kiyosu 清洲 Castle Town site (fifteenth and sixteenth centuries), Aichi; Standard Chronology D (512-1322) derived from artifacts discovered at the Toba 鳥羽 Detached Palace (eleventh and twelfth centuries) site, Kyoto and those at the Kusado Sengen-chō 草戸千軒町(Medieval Age) site, Hiroshima; Standard Chronology E (37, B.C.-A.D.838) derived from artifacts discovered in the Heijō 平城 Capital (710-784) site, Nara; and Standard Chronology F (317, B.C.-A.D. 258) derived from artifacts discovered at the Yayoi (fourth century, B.C. to A.D. third century) and Kofun (fourth to sixth centuries) sites.

We also built chronologies derived from *sugi* cedar trees, but none of them overlapped with one another. They were: Standard Chronology A (1779-1986) taken from a modern tree; Standard Chronology B (405-1285) taken from artifacts discovered at sites in the Tohoku District and cross-correlated with the Index Master Chronology of cypress; and Standard Chronology C (420, B.C.-A.D.265) derived from artifacts discovered at sites in the Tokai region (the Pacific Coast of the Chubu District) and cross-correlated with the Standard Chronology F of cypress.

In addition, we built two Chronologies utilizing *kōyamaki* pine trees. A 556 year ring pattern derived from artifacts discovered in the Heijō Capital cross-correlated with the Standard Chronology E of cypress and spanned from A.D.186 to 741. In the same way, an artifact discovered at the Shijō Kofun 四条古墳 burial site in Nara gave rise to a chronology between 286 and 695.

We then applied these chronologies to artifacts, early architecture, and crafts in an attempt to date them. Here are some examples: the Shijō Kofun, Nara, the Seta no Kara Hashi 瀬田唐橋 Bridge site in Shiga Prefecture, the Miyamachi 宮町 site, Shiga Prefecture, and the Hotta no Saku 扠田柵(Hotta Fort) site in the present Iwate Prefecture.

At the Shijō Kofun, 448 was the latest dendrochronological date obtained from numerous artifacts made of *kōyamaki* pine, which had presumably been used for mortuary rituals, and were discovered in the moat surrounding the Kofun. In other words, the Kofun was built sometime after 448. Since the absolute dating of *kofun* or burial mounds built between the late third and sixth centuries had always been debated, we believe that dendrochronology would be a useful method by which to determine absolute dates.

We applied dendrochronology to try to estimate when a bridge was built for the first time over the Seta River. Archaeologists excavated struc-

tures of a bridge near the present Seta no Kara Hashi Bridge 3.5 meters below the bottom of the River. We utilized several long pieces of timber and lumber of 20 to 50 centimeters in diameter or in length. We confirmed that the bridge was constructed after 607, and consider it probable that this was the strategic bridge the control over which was an important aspect of a civil war in 672.

A ring pattern taken from the lower portion of a wooden pillar discovered at the Miyamachi site cross-matched with the Standard Chronology E of cypress, and we could confirm that the tree for the pillar was cut down between 742 and the beginning of 744. This made it very highly likely that the site was a part of the Shigaraki 紫香樂 Palace which was occupied from the eighth month of 742 (on the lunar calendar) and the fifth month of 745.

In the same way, we could determine that the construction of the Hotta no Saku 弘田柵 was begun between 801 and 802. There was a cross-matching of ring patterns yielded from logs of cedar (a square of approximately 30 centimeters in length) which surrounded the site and the Standard Chronology B of cypress. This refuted the hypothesis that the site was the historically known Okachi 雄勝 Castle built in 759.

Our study of dendrochronology made considerable contributions to architectural history and art history as well. Examples of dendrochronological applications included Hōryū-ji, a Nara Period horse painting, a Kamakura Period guardian figure at Tōdai-ji 東大寺, Buddha figures in Yamaguchi Prefecture, Ha'ushiwake 波宇志別 Shrine in Akita Prefecture, Wakamiya Hachiman 若宮八幡 Shrine in Nagano Prefecture and a Medieval wooden bent-box in Kyoto:

We applied dendrochronology to date the western complex of the Hōryū-ji. There had been two major hypotheses concerning the dating of Hōryū-ji: a) the temple was built around 610 during the reign of Empress Suiko 推古; and b) the temple was burnt down around 670 and rebuilt at the end of the seventh or the beginning of the eighth century. The number of rings of the central pillar of the pagoda in the western complex was 351, corresponding to the years 241 to 591. The pillar had been extensively modified from the original tree trunk by largely removing the exterior portion and the central portion; it would be unlikely that only the outermost twenty rings were removed for this modification. We found it very unlikely that the pagoda was built during the reign of Empress Suiko around 610.

Next, we were able to give a more specific date to a small painting. In 1989 we discovered in a garbage pit in the Heijō Capital a paint-

ing of a horse done on a cypress board of 27 by 19.5 centimeters and 0.7 centimeters in thickness. Since this board still kept the most exterior portion of the original tree, we could date the outer most ring to the year 728. We could also date the pit to 737 by a wooden tablet with a dated inscription. The horse was probably painted between 728 and 737. This dating was felicitous because the horse was one of the few surviving examples of the eighth century paintings in Japan.

Next, our ability to achieve this degree of specificity in dating, we could confirm a historical record. In 1181, one year after Tōdai-ji was burnt down in a civil war, priest Chōgen 重源 and his collaborators began to reconstruct the temple. Two gigantic guardian figures (4.5 meters in height), housed in the south gate of the temple, were built between the seventh month and eleventh month of 1203 (on the lunar calendar). In 1989, there was an overhaul of one of the figures, so we took the opportunity to apply dendrochronological dating to the parts of the figure which still retained bark. Original trees utilized for this statue were cut down in the winter of 1196, 1199, and 1201 or during the following spring of those years. Further, we discovered that one of the ring patterns taken from the statue was almost identical to that of a Buddha figure in the present Yamaguchi Prefecture—approximately 400 kilometers apart. Since it was historically known that Chōgen brought timber for structural parts of the temple, it was quite likely that wood for the guardian figures also came from Yamaguchi.

In other cases, finally, dendrochronological dates could contradict dates determined according to stylistic evolution. In and the vicinity of the Tokuchi 徳地 Town, Yamaguchi Prefecture, there were many Buddha figures in various temples, which were stylistically dated to the late twelfth century. We applied dendrochronology to fifteen of them and discovered that the most recent rings of trees used in six of the figures were formed in the 1190s. Since the exterior portion, and in some cases the central portion of trees, had largely been removed, the trees must have been cut down in the *thirteenth* century. The earlier dating should be reconsidered.

Structural parts of the Kagura-den 神楽殿 (Hall of Ritual Music and Dance) at the Ha'ushiwake Shrine yielded dendrochronological dates of 1177 and 1195 by cross-matching with the Standard Chronology B of cedar. Since both of the structural parts kept the bark, the original trees were cut down around that time in the twelfth century. According to stylistic criteria, the Kagura-den had been estimated to date from the fifteenth century. When these Shrine structures are overhauled sometime in the future, we should more thoroughly investigate the dates of the structures by apply-

ing dendrochronology.

Similarly, major pillars of the Hon-den 本殿(Main Hall) at the Wakamiya Hachiman Shrine were dendrochronologically dated to be after 1614, the structure was stylistically considered to be of the Momoyama Period (1582-1600).

Further, this method was effective in detecting forgeries. A certain temple in Kyoto Prefecture owned a wooden bent-box of cypress (a cylindrical container) which had the inscription dated to 1233 on the bottom. We discovered, however, that the most recently formed tree-ring was dated to 1576. The container was a forgery !

In addition to architectural structures, art objects, and artifacts, dendrochronology is effective in dating natural disasters. A standing *hinoki* cypress of 1.4 meters in diameter with the bark completely preserved was discovered underground in the Susono 裾野 City at the foot of Mt. Fuji 富士. Taking the topography of the vicinity into consideration, it was probable that lava issuing from an eruption of Mt. Fuji dammed up a river, thereby submerging the whole forest and killing the trees. If this was indeed the case, we could estimate the date of an eruption of Mt. Fuji. By cross-matching with the Standard Chronology E of cypress, we could date the standing tree to A.D.833.

Similar discoveries of buried forests which might indicate natural disasters were reported from all over Japan. In the Mamurogawa 真室川 Town, Yamagata Prefecture, some twenty cedar trees were similarly discovered six meters below the ground. By cross-matching with the Standard Chronology B of cedar, we confirmed that the most recent ring was formed in A.D.850. In other words, a natural disaster which buried a forest hit this area between the winter of 850 and spring of 851.

In Futatsu'i ニツ井 Town, Akita Prefecture where the people believed in a tale that several thousand *koku*² of wild cedars were buried by a landslide during the Tokugawa Period (1600-1868), numerous large trunks of cedars and Japanese oaks of approximately 1.5 meters in diameter were discovered a few meters below the ground. We dated 754 rings of a cedar tree by cross-matching with the Standard Chronology A of cedar and discovered that the most exterior ring was formed in A.D.958. The forest was buried in the middle of the tenth century, much earlier than the people believed.

2 a unit of volume used widely in Japan before the Meiji Restoration of 1898; in the case of trees, one *koku* equaled to 0.278 cubic meters.

In the southern part of Akita Prefecture, there was an area where a large forest was buried more than ten meters below the ground. Most of the trees were cedar and some exceeded four meters in diameter. Taking into consideration the soil of the area and the way in which the trees had fallen, we hypothesized that an enormous mud flow resulted from an eruption of Mt. Chōkai 鳥海 some ten kilometers southeast of the forest. Radiocarbon dates showed that the disaster happened about 2600 B.P. We applied dendrochronology to eight samples taken from the buried forest, six of which were with barks completely preserved. The most recent rings were formed at the same time; it was likely that one eruption caused this complete destruction. The fact that the most recent rings contained fully mature cells indicated that the eruption occurred sometime between the winter of a certain year and the spring of the following year. We derived a ring pattern which spanned 848 years, but it remained as a floating chronology; it was too old to be cross-matched with any standard chronologies. Once we extend our Index Master Chronology to 2600 B.P., we could confirm the precise date of this disaster.

In addition to the dendrochronological advances discussed above, we progressed in the application of dendroclimatology. The ring pattern we used was derived from thirty samples of living cypress trees in Nagano Prefecture and the Standard Chronology B which was presumably based on timber produced in the same prefecture. We also collected the data concerning the number of rainy days from April to September in each year for the past 170 years from a local weather station and from diaries kept by local people before the weather station started. We filled missing data (years for which we did not have records of rainy days) by a one-dimensional autoregressive process. We discovered that our reconstructed 170 year-long data of rainy days per annum³ closely corresponded to the current precipitation data in Nagano Prefecture. We concluded that the number of rainy days was appropriate for climatic data.

In order to determine the most influential climatic factor in tree growth, we tried correlating the number of rainy days and average temperature to tree ring width. Tree grew fastest when there were many rainy days temperature did not rise.

We then compared the power spectra of tree ring width and of the per annum rainy days. We discovered cycles of approximately six years, eight

3 Throughout this section, by "per annum rainy days" we mean the total number of rainy days from April to September. Such data sets were chosen as the climatic data because tree rings grow most from spring to fall.

years, and twenty-three years to be common to both sets of data. This indicated to us a very good correlation between the tree ring width and per annum rainy days.

Owing to this correlation, we proposed a system in which climatic data represented by per annum rainy days were input and tree ring width data were output, by the process of an autoregressive moving average function. In order to identify per annum rainy days from the tree ring width data, we developed a correlation function. The most effective function for this purpose was a process with three autoregressive and two moving average coefficients. Using the identified parameters of this process, we reconstructed the per annum rainy day for eighty years beginning in 1760. We derived the relationship between the tree ring width data and per annum rainy days from 1009 to 1984, assuming the linear system we had used previously between the two factors. We standardized the primary tree ring width data for these 976 years by the spline function. We used the Kalman filtering algorithm, including the U-D observation updated theorem, to identify the system parameters. In the process of parameter identification, every parameter gradually converged to a certain value.

By utilizing these system parameters, we calculated per annum rainy days for the 170 years for which we originally had good weather records. Our calculation was in basic agreement with the actual data. In the same way, we could reconstruct the per annum rainy days for approximately 800 years before 1813. These results of our research showed that the application of dendroclimatology was possible in Japan.

Another major contribution of our research was the applicability of dendrochronology and dendroclimatology to Palaeolithic samples. At the Tomizawa 富沢 site, Sendai City, Miyagi Prefecture, excavators discovered features three meters below the ground, which suggested that the inhabitants had been manufacturing stone tools at a camp fire. We could build a ring pattern spanning 323 years, derived from cross-correlation of three samples of genus *Larix* (a kind of larch). Since the site was dated by carbon 14 to have been more than 20,000 years old, we could not dendrochronologically determine the date of the site, as our Index Master Chronology did not extend beyond 317, B.C. Nonetheless, this result showed the potential application of tree ring studies to the Palaeolithic Period.

In summary, our ten year project (including the preliminary research started in 1979) has shown that the application of dendrochronology and dendroclimatology to Japanese trees is possible. Our Index Master Chronolo-

gy of cypress extends back to 317,B.C. The Master Chronology of cedar of the Tohoku District extends to A.D.405, and that of cedar of the Pacific Coast of the Chubu District to 420,B.C. We have certainly progressed to the stage of application to other disciplines such as archaeology, architectural history, art history and the history of natural disasters in Japan.